

# On the interface between lattice results and $\chi$ PT

H. Leutwyler  
University of Bern

Colloquium in memory of Jan Stern

FROM CURRENT ALGEBRA TO THE STANDARD MODEL AND BEYOND

October 2, 2009, Institut Henri Poincaré, Paris



Bern, 1978

I had a wonderful time working together with Jan

For reminiscences, see talk given at IN2P3, Orsay, June 2008: [www.leutwyler.itp.unibe.ch](http://www.leutwyler.itp.unibe.ch)

# Standard Model at low energies

- Low energies ( $E \ll M_W$ ): weak interaction is frozen
- ⇒ Standard Model reduces to QCD + QED
- Lagrangian only involves  $g_s, \theta, e$ , fermion masses
- ⇒ Precision theory for cold matter ( $T \ll M_W$ ), size and structure of atoms, solids, etc.
- QED is infrared stable, characterized by pure number, which happens to be small,  $1/137$
- ⇒ QED can be accounted for with perturbation theory
- Hadrons at low energies: SM = QCD + corrections

## Pièce de résistance: QCD

- $\exists$  many models that resemble QCD: instantons, monopoles, bags, superconductivity, gluonic strings, linear  $\sigma$  model, hidden gauge, NJL, AdS/CFT, but ...
- Nonperturbative methods needed  
⇒ Progress in understanding is slow
- Model independent methods:
  - Numerical simulation on a lattice
  - Sum rules, dispersion relations
  - Effective field theory ( $\chi$ PT)

# Hidden symmetries in particle physics

Already in 1960, Nambu realized that

1.  $SU(2)_L \times SU(2)_R$  is an approximate symmetry of the strong interaction
2. The symmetry is "hidden", "spontaneously broken":  $|0\rangle$  invariant only under the isospin subgroup  $SU(2)_{L+R}$
3. The spontaneous breakdown of an exact symmetry entails massless particles
4. For the strong interaction, the pions play this role
5. The pions are not massless, only light, because the symmetry is only an approximate one

Nobel Prize 2008

Explains why the energy gap of the strong interaction is so small :  $M_\pi \simeq 135 \text{ MeV}$

When Nambu proposed this idea, the origin of the symmetry was mysterious

Approximate symmetries ? Partially conserved currents ?

For gauge theories like QCD, approximate symmetries do occur naturally

# Chiral symmetry

Where is Nambu's hidden approximate symmetry in QCD ?

- QCD with  $N_f$  massless quarks: Lagrangian has an exact chiral symmetry:  $SU(N_f)_L \times SU(N_f)_R$
- $|0\rangle$  is symmetric only under the subgroup  $SU(N_f)_{L+R}$   
Symmetry is spontaneously broken
- ⇒ Spectrum contains  $N_f^2 - 1$  Nambu-Goldstone bosons
- $m_u$  and  $m_d$  happen to be small
- ⇒  $SU(2)_L \times SU(2)_R$  is an approximate symmetry of QCD
  - broken spontaneously
  - ⇒  $|0\rangle$  not invariant
  - broken explicitly by mass term  $m_u \bar{u}u + m_d \bar{d}d$
  - ⇒  $\mathcal{L}_{\text{QCD}}$  not invariant
  - $m_u, m_d$  are very small  $\rightarrow$  symmetry is nearly exact

# Chiral perturbation series

- For  $m_u = m_d = 0$ , pion exchange gives rise to poles and branch points at  $p = 0$ 
  - ⇒ Low energy expansion is not a Taylor series, contains infrared singularities
- Properties of the Nambu-Goldstone bosons are governed by the hidden symmetry that is responsible for their occurrence
- NG bosons of low momentum interact only weakly: can treat the momenta as well as  $m_u, m_d$  as perturbations
  - ⇒ Chiral perturbation series: simultaneous expansion of the matrix elements in powers of  $p, m_u, m_d$

# Effective Lagrangian

- Formulation in terms of an effective Lagrangian

Weinberg 1967, Coleman, Wess, Zumino, Callan, Dashen, Weinstein 1969

- Lagrangian  $\supset$  massless Nambu-Goldstone Bosons

- ⇒ Perturbation series has infrared singularities

Li + Pagels 1971, Langacker + Pagels 1973

Weinberg 1979, Gasser + Zepeda 1980, Gasser 1981

Singularities due to NG bosons can be worked out with an effective field theory  
“Chiral Perturbation Theory”

- $\chi$ PT reproduces the low energy structure of QCD, order by order in the expansion in powers of  $p$ ,  $m_u$ ,  $m_d$



# Plethora of low energy constants

- $\chi$ PT merely exploits the symmetries of QCD: yields the general solution of the Ward-Takahashi identities
- $\mathcal{L}_{eff}$  contains all functions that can be formed with the pion field and its derivatives, only subject to the condition that the sum is chirally invariant
- Order in number of derivatives (powers of momentum)
- ⇒ Number of terms in  $\mathcal{L}_{eff}$  rapidly grows with the order:  
LO: 2, NLO: 7, NNLO: 53, ...
- Symmetries only relate – do not determine
- In principle, the effective theory is exact:  
yields expansion of QCD Green functions in  $p, m_q$

# Illustration: energy gap of QCD

- Energy gap of QCD:  $M_\pi$
- Ignore e.m. self energy,  $e = 0$ , pure QCD
- ⇒  $M_\pi$  is a function of  $\Lambda_{\text{QCD}}, m_u, m_d, \dots, m_t$
- How does  $M_\pi$  depend on  $m_u, m_d$  ?

Chiral symmetry:  $M_\pi \rightarrow 0$  for  $m_u, m_d \rightarrow 0$

- Leading order formula (tree level of  $\chi$ PT):

$$M_\pi^2 = (m_u + m_d)B$$


Gell-Mann, Oakes, Renner 1968

- The coefficient is determined by the quark condensate:


$$B = \frac{|\langle 0 | \bar{u}u | 0 \rangle|}{F_\pi^2}$$

# Lattice results for $M_\pi$

- GMOR formula can now be checked on the lattice: determine  $M_\pi$  as a function of  $m_u = m_d = m$



Lüscher, Lattice conference 2005



ETM collaboration, hep-lat/0701012

These plots are for QCD with  $N_f = 2$ . Behaviour in full QCD looks very similar.

# Lattice

- Quality of data is impressive
- No quenching, quark masses are sufficiently light
- ⇒ Legitimate to use  $\chi$ PT for the extrapolation to the physical values of  $m_u, m_d$

- Proportionality of  $M_\pi^2$  to

$$m_{ud} \equiv \frac{1}{2}(m_u + m_d)$$

holds out to  $m_{ud} \simeq 10 \times m_{ud}^{\text{phys}}$

- Main limitation: systematic uncertainties from lattice artifacts, continuum extrapolation, finite size effects, etc.

## Expansion of $M_\pi^2$ in powers of $m_u, m_d$

- GMOR formula represents leading term of  $\chi$ PT
- Correction of first nonleading order:

$$M_\pi^2 = M^2 \left\{ 1 - \frac{M^2}{32\pi^2 F_\pi^2} \bar{\ell}_3 + O(M^4) \right\}$$

$$M^2 \equiv B(m_u + m_d)$$

$\ell_3 \in \mathcal{L}_{\text{eff}}$  depends logarithmically on running scale

- What counts is the running coupling at scale  $M_\pi$ :

$$\bar{\ell}_3 = \ell n \frac{\Lambda_3^2}{M_\pi^2}$$

⇒ Expansion of  $M_\pi$  contains a chiral logarithm


Langacker + Pagels 1973, Gasser + Zepeda 1980, Gasser 1981

# Size of the low energy constant $\bar{\ell}_3$

Crude estimate, based on  $SU(3)_L \times SU(3)_R$ :  $\bar{\ell}_3 = 2.9 \pm 2.4$

Gasser & L. 1984

Lattice allows more accurate determination:



$$\bar{\ell}_3 = \ln \frac{\Lambda_3^2}{M_\pi^2}$$

Result of RBC/UKQCD 08, for instance:  $\bar{\ell}_3 = 3.13 \pm 0.33 \pm 0.24$   
*stat* *sys*

## Expansion of $F_\pi$ in powers of the quark mass

- Also contains a logarithm at NLO:

$$F_\pi = F \left\{ 1 + \frac{M^2}{16\pi^2 F^2} \ell n \frac{\Lambda_4^2}{M^2} + O(M^4) \right\}$$

$$M_\pi^2 = M^2 \left\{ 1 - \frac{M^2}{32\pi^2 F^2} \ell n \frac{\Lambda_3^2}{M^2} + O(M^4) \right\}$$

$F$  is value of pion decay constant in limit  $m_u, m_d \rightarrow 0$

- Structure is the same, coefficients and scale of logarithm are different
- Quark mass dependence of  $F_\pi$  can also be measured on the lattice  $\Rightarrow$  measurement of  $\Lambda_4$
- Alternative method: determine the scalar form factor of the pion, radius  $\langle r^2 \rangle_s \leftrightarrow \bar{\ell}_4 = \ell n(\Lambda_4^2/M_\pi^2)$

Colangelo, Gasser & L. 2001

## Lattice determination of scalar radius

- Scalar form factor can be measured on the lattice
- Most recent lattice determination:

$$\langle r^2 \rangle_s = 0.617 \pm 0.079_{\text{stat}} \pm 0.066_{\text{syst}} \text{ fm}^2$$

JLQCD/TWQCD collaboration, arXiv:0905.2465

This is consistent with the value obtained on the basis of dispersion theory,


$$\langle r^2 \rangle_s = 0.61 \pm 0.04 \text{ fm}^2$$

Colangelo, Gasser & L. 2001

but the uncertainties in the lattice result are still large



# Size of $\ell_4$



$$\bar{\ell}_4 = \ln \frac{\Lambda_4^2}{M_\pi^2}$$

Lattice results are consistent with value obtained from dispersion theory, uncertainties are comparable

## $\pi\pi$ interaction

- Symmetry fixes the interaction among the Nambu-Goldstone bosons

- LO formulae for the S-wave scattering lengths:

$$a_0^0 = \frac{7M_\pi^2}{32\pi F_\pi^2}, \quad a_0^2 = -\frac{M_\pi^2}{16\pi F_\pi^2} \quad \text{Weinberg 1966}$$

- NLO corrections are determined by  $\ell_3, \ell_4$  Gasser + L. 1983

- $\pi\pi$  scattering amplitude known to NNLO

Bijnens, Colangelo, Ecker, Gasser + Sainio 1996


- Uncertainty in predictions for  $a_0^0, a_0^2$  is dominated by the uncertainty in the low energy constants  $\ell_3, \ell_4$

⇒ Can make use of the lattice results for these

- Contributions from higher order couplings are tiny

Guo + Sanz-Cillero arXiv:0904.4178

# Consequence of lattice results for $l_3, l_4$



The plot represents beautiful physics: experiment as well as theory  $\Rightarrow$  talk by Jürg Gasser

## Extension to $SU(3)_L \times SU(3)_R$


- In the theoretical limiting case  $m_u = m_d = m_s = 0$  QCD acquires an exact  $SU(3)_L \times SU(3)_R$  symmetry

Is  $m_s$  small enough for this to represent a useful approximate symmetry ?

- Theoretical reasoning
  - $SU(3)_{L+R}$  (eightfold way) is an approximate symmetry
  - Typical size of  $SU(3)_{L+R}$  breaking:  $\frac{F_K}{F_\pi} = 1.19 \pm 0.01$
  - Only coherent way to understand this in QCD:  
The mass differences  $m_s - m_d$ ,  $m_d - m_u$  must be small, can be treated as perturbations
  - Since  $m_u, m_d \ll m_s$
- ⇒  $m_s$  is small,  $SU(3)_L \times SU(3)_R$  must be an approximate symmetry, breaking not larger than for  $SU(3)_{L+R}$

## Expansion in powers of $m_u, m_d, m_s$

- Expansion in powers of  $m_u, m_d, m_s$  ought to work, but expect convergence to be comparatively slow
- Lattice results:  $M_\pi^2 \propto m_{ud}$  holds out to  $10 \times m_{ud}^{\text{phys}}$
- $m_s$  is larger than that:  $m_s \simeq 27 \times m_{ud}$



Compare


$$\frac{F_K}{F_\pi} \simeq 1.19$$

# Three light quarks: interface between lattice and $\chi$ PT

- Steady progress in simulating QCD with light quarks, but the quark masses used are still too large for the NLO formulae of  $\chi$ PT to work
- $M_\pi$  OK, but  $M_K$  too large
- Three options
  - Use smaller quark masses
  - Extrapolate only in  $m_u, m_d$ , keep  $m_s$  fixed
  - Account for NNLO contributions
- Some lattice analyses do allow for NNLO contributions, but the chiral logarithms are accounted for only to NLO
- $\exists$  discrepancies between different lattice results  
In part, these may arise from nonperturbative renormalization effects  
Some of the collaborations still use perturbative renormalization

$\Rightarrow$  Illustrate this with the results for  $m_s$

# Mass of the strange quark



Lattice and sum rule results agree within errors  
 Can expect significant progress in lattice determinations very soon

## Relative size of $m_u, m_d, m_s$

- $M_{\pi^+}^2 = (m_u + m_d) B_0 + O(m^2)$   
 $M_{K^+}^2 = (m_u + m_s) B_0 + O(m^2)$   
 $M_{K^0}^2 = (m_d + m_s) B_0 + O(m^2)$
- $\chi$ PT relates  $B_0$  to the quark condensate, but does not predict its size  $\Rightarrow$  no prediction for size of quark masses

- Account for e.m. self energies at tree level of  $\chi$ PT and drop effects of second order in isospin breaking

$$\frac{m_u}{m_d} = \frac{M_{K^+}^2 - M_{K^0}^2 + 2M_{\pi^0}^2 - M_{\pi^+}^2}{M_{K^0}^2 - M_{K^+}^2 + M_{\pi^+}^2} = 0.56$$

$$\frac{m_s}{m_d} = \frac{M_{K^+}^2 + M_{K^0}^2 - M_{\pi^+}^2}{M_{K^0}^2 - M_{K^+}^2 + M_{\pi^+}^2} = 20.2$$

Weinberg 1977

- Corrections from higher orders ? Could they strongly modify the numerical values ?  $m_u = 0$  ?



## Higher orders

- $\nexists$  scalar probe analogous to  $\gamma$ ,  $W^\pm$
- ⇒ Quark masses cannot be determined from phenomenology alone, not even their ratios

Kaplan & Manohar 1986

- At LO,  $\chi$ PT does determine the quark mass ratios
- At NLO, there is only one relation without unknowns:

$$Q^2 \equiv \frac{m_s^2 - m_{ud}^2}{m_d^2 - m_u^2} = \frac{M_K^2 - M_\pi^2}{M_{K^0}^2 - M_{K^+}^2} \frac{M_K^2}{M_\pi^2} + \text{NNLO} + \text{e.m.}$$

$M_K, M_\pi$ : mean masses of the two multiplets

Gasser & L. 1985

The relation correlates the two ratios

Value of  $Q \rightarrow$  ellipse in the plane  $\left(\frac{m_u}{m_d}, \frac{m_s}{m_d}\right)$

- Weinberg's leading order formulae give  $Q = 24.3$ .

$$\eta \rightarrow \pi^+ \pi^- \pi^0$$

- Critical input for value of  $Q$  is the "Dashen theorem": e.m. self energies are accounted for only at tree level

- $\eta$  decay allows an independent determination of  $Q$

Gasser & L. 1985

- Dispersive analysis of the decay amplitude

Kambor, Wiesendanger & Wyler 1996, Anisovich & L. 1996, Walker 1998

- In  $\eta \rightarrow 3\pi$ , the e.m. contributions are suppressed

Bell & Sutherland 1968

- ⇒ Uncertainties are smaller


Quantitative analysis of e.m. contributions: Ditsche, Kubis & Meißner 2009

- Update of Walker's calculation with the current experimental information ⇒  $Q = 22.4 \pm 0.8$ , to be compared with  $Q = 24.3$  from Dashen theorem

- Comprehensive analysis of  $\eta \rightarrow 3\pi$  is under way

PhD thesis of Stefan Lanz, in preparation

# $\chi$ PT at leading and first nonleading order



## Where on the ellipse ? $m_u = 0$ ?

- The vacuum angle  $\theta$  breaks CP
- Chiral symmetry ensures that  $\theta$  can enter physical quantities only via  $\det \mathcal{M} \times e^{i\theta}$
- If  $m_u$  is zero  $\rightarrow \det \mathcal{M}$  vanishes  $\rightarrow \theta$  without physical significance  $\rightarrow$  QCD invariant under CP
- Quite a few authors advocated  $m_u = 0$  as the solution of the strong CP problem, possibly some still do ...

Nice idea, but amounts to trading one puzzle for the other:

- If  $m_u$  were zero, the Weinberg formula for  $m_u/m_d$  would turn into a prediction for  $M_{K^0} - M_{K^+}$ :

$$M_{K^0} - M_{K^+} = \frac{2M_{\pi^0}^2 - M_{\pi^+}^2}{M_{K^0} + M_{K^+}} + \text{NLO}$$

$\uparrow$

3.9 MeV

experiment

$\uparrow$


16.9 MeV

theory

$$m_u = 0 ?$$


- If  $m_u$  were zero, then  $\chi$ PT would be in conflict with the observed mass pattern of the NG bosons
  - chiral series could not be truncated at low orders
  - $SU(3)_L \times SU(3)_R$  not an approximate symmetry
  - Success of Gell-Mann-Okubo formula accidental etc.
- Leading order formula for  $M_{K^0} - M_{K^+}$  is off by less than a factor of 2 only if  $0.7 > m_u/m_d > 0.33$

# Allowed range of mass ratios



- All of the lattice results are in the range allowed by  $\chi$ PT  
None is consistent with the solution  $m_u = 0$  of the strong CP problem
- The MILC collaboration rules this solution out at  $10\sigma$
- ⇒ Nature solves the strong CP problem differently

# Results for quark mass ratios



# Paramagnetic inequalities

- Consider the limit  $m_u, m_d \rightarrow 0$ ,  $m_s$  physical
  - $F$  is the value of  $F_\pi$  in this limit
  - $\Sigma$  is the value of  $|\langle 0 | \bar{u}u | 0 \rangle|$  in this limit
  - $F_0, \Sigma_0$  are the values of  $F, \Sigma$  in the limit  $m_s \rightarrow 0$
- Inequalities set up by Jan and collaborators:  
both  $F$  and  $\Sigma$  should decrease if  $m_s$  is taken smaller

$$F > F_0, \Sigma > \Sigma_0$$

for a recent discussion, see S. Descotes-Genon at Lattice 2007


- If  $N_c$  is large  $\rightarrow F$  and  $\Sigma$  become independent of  $m_s$
- $\Rightarrow F/F_0 - 1$  and  $\Sigma/\Sigma_0 - 1$  violate the OZI rule
- The lattice results confirm the parametric inequalities, but do not yet allow to draw conclusions about the size of the OZI-violations



# Quark condensate

$$\Sigma = |\langle 0 | \bar{u}u | 0 \rangle|_{m_u, m_d \rightarrow 0}$$

$$\Sigma_0 = |\langle 0 | \bar{u}u | 0 \rangle|_{m_u, m_d, m_s \rightarrow 0}$$



Central values of RBC/UKQCD and PACS-CS for  $\Sigma/\Sigma_0$  lead to qualitatively different conclusions concerning OZI-violations  $\Rightarrow$  discrepancy indicates large systematic errors

## Lattice determination of $V_{us}$ , $V_{ud}$

- Rely on Standard Model, where

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$$

- Precision data on  $K$ -decays imply



$$|V_{us}|f_+(0) = 0.21661(47)$$

$$\left| \frac{V_{us}F_K}{V_{ud}F_\pi} \right| = 0.27599(59)$$

- ⇒ Since  $V_{ub}$  is tiny and known to good accuracy,  $V_{ud}$ ,  $f_+(0)$ ,  $F_K/F_\pi$  are all determined by  $V_{us}$


- Lattice allows two independent ways to measure  $V_{us}$ :  
calculate  $f_+(0)$  or calculate  $F_K/F_\pi$

# Lattice results for $f_+(0)$ and $F_K/F_\pi$



- FLAG estimate combines the lattice data for  $f_+(0)$  with those for  $F_K/F_\pi$

# Lattice results for $V_{us}$ and $V_{ud}$



FLAG preliminary

- Confirms nuclear  $\beta$  decay value for  $V_{ud}$  within errors
- $\tau$  decay: physics beyond the Standard Model ?

# Trying to understand the size of the low energy constants

- $SU(2)_L \times SU(2)_R$ : can understand the size of all NLO couplings in terms of resonance exchange Gasser + L. 1984
- Also true for  $SU(3)_L \times SU(3)_R$  Ecker, Gasser, Pich, de Rafael 1989
- $\chi$ PT formulae have been worked out to NNLO for many quantities of physical interest Bijnens and collaborators
- Formulae involve new unknown low energy constants
- "Resonance Chiral Theory": couplings of higher order, effective Lagrangian for e.m. + weak interactions ...  
Gonzalez-Alonso, Guo, Pich, Portoles, Prades, Rosell, Ruiz-Femenia, Sanz-Cillero ...
- Comprehensive review of current state of the art:  
Bijnens, arXiv:0904.3713 (Valencia 2009)

# Problem with Resonance Chiral Theory in case of $f_+(0)$

- Form factor known to NNLO

Post + Schilcher 2002, Bijens + Talavera 2003

- Account for isospin breaking, use  $R_\chi$ PT estimates for the low energy constants

$$\Rightarrow f_+(0) = 0.986(7)$$

Kastner + Neufeld 2008


- To be compared with

$$0.961(5) (\beta \text{ decay}) \text{ or } 0.962(5) (\text{lattice})$$

(These numbers are obtained by assuming that the CKM matrix is unitary)

Discrepancy amounts to  $2.9 \sigma$  and  $2.8 \sigma$ , respectively

# Compare with lattice results



- $1 - f_+(0)$  is a symmetry breaking effect
- No problem at NLO: parameter free prediction
- ⇒  $R_{\chi PT}$  does not appear to account properly for the symmetry breaking effects at NNLO

## Problems with scalar meson dominance ?

- Quark mass term in  $\mathcal{L}_{\text{QCD}}$  is a scalar operator
- Matrix elements dominated by scalar resonances ?  
Can the *dependence on the quark masses* be accounted for with scalar meson dominance ?
- Rapidly rising  $\pi\pi$  continuum (large chiral logs),  $\sigma$  makes a broad bump, narrow peak from  $f_0(980)$ , glueballs, etc.
- Failure of scalar meson dominance may be the origin of the problem

more detailed discussion in Erice lectures 2007



# Conclusions

# Conclusions

- Expansion in powers of  $m_u, m_d$  yields a very accurate low energy representation of QCD
- Lattice yields remarkably coherent and significant results for pion physics already now

Low energy pion physics is a precision laboratory  
Theoretical tools:  $\chi$ PT, lattice, dispersion theory

- Limitations:
  - Low energies
  - e.m. interaction must properly be accounted for
  - Calculations cannot be done on back of an envelope

# Conclusions

- $m_u \neq 0$

Nature solves the strong CP problem differently

- $m_s = 98 \pm 11 \text{ MeV}$  FLAG 09 (preliminary result)  
MS scheme, scale 2 GeV

Lattice results confirm sum rule estimates within errors

- For the physical values of  $m_u$ ,  $m_d$ ,  $m_s$ , the leading order terms in the chiral perturbation series of  $M_\pi$ ,  $M_K$ ,  $F_\pi$ ,  $F_K$  do represent a decent approximation

- Summary of current knowledge of quark mass ratios:

$$\frac{m_u}{m_d} = 0.47 \pm 0.08$$

$$\frac{m_s}{m_d} = 19.7 \pm 1.5$$

to be compared with Weinberg's LO formulae, which give 0.56 & 20.2, respectively

# Conclusions

- Lattice results indicate that the NLO contributions in  $M_\pi, M_K, F_\pi, F_K$  do dominate the corrections
- ⇒  $\chi$ PT does appear to work for  $SU(3)_L \times SU(3)_R$  as well
- Extension to kaon physics is making progress
  - Except for a few selected quantities, kaon physics is still at an exploratory stage
  - Representations of many quantities of interest are available to NNLO of  $\chi$ PT ⇒ Bijnens et al.
  - Main problem at NNLO: the current knowledge of the LECs is rudimentary
  - The  $R\chi$ PT estimates for  $f_+(0)$  illustrate the problem
  - There was a problem with the  $R\chi$ PT estimates also for  $K \rightarrow \pi\pi$ , but this puzzle appears to be solved

Cirigliano, Ecker + Pich, Phys. Lett. 2009

# Conclusions

- Many open issues:
  - $M_K = 600$  MeV is beyond reach of  $\chi$ PT
  - Better determination of some of the LECs needed
  - In particular, a meaningful comparison of many of the  $\chi$ PT results in kaon physics with experiment requires better knowledge of those LECs that determine the dependence on the quark masses
  - Size of Okubo-Iizuka-Zweig rule violations ?
  - e.m. self energies, corrections to Dashen Theorem ?

Significant progress at the interface between lattice and effective field theory methods is ante portas

# Spares

## Phase of final state in $K \rightarrow \pi\pi$

- $K \rightarrow \pi\pi$  decay: value of  $\delta_0^0 - \delta_0^2$  at  $s = M_K^2$ 
  - $\pi\pi$  phase shifts accurately known from dispersion theory  $\delta_0^0 - \delta_0^2 = 47.5^\circ \pm 1.5^\circ$  Colangelo, Gasser, L. 2001
  - In the determination from  $K \rightarrow \pi\pi$  via Watson theorem, isospin breaking is enhanced because of the  $\Delta I = \frac{1}{2}$  rule
  - Complete analysis to NLO Cirigliano, Ecker, Neufeld + Pich 2004
  - Recent update of the numerics yields

$$\delta_0^0 - \delta_0^2 = 52.5^\circ \pm 0.8^\circ_{\text{exp}} \pm 2.8^\circ_{\text{th}}$$

Cirigliano, Ecker + Pich, Phys. Lett. 2009

- Remaining difference amounts to  $1.5 \sigma$

# Large $N_c$

- In the large  $N_c$  limit, the  $\eta'$  also becomes a Nambu-Goldstone boson
- ⇒ Can extend  $\chi$ PT to include the  $\eta'$ , systematic expansion in powers of  $m_u$ ,  $m_d$ ,  $m_s$  and  $1/N_c$
- In this framework, there is no ambiguity at NLO
- Triangle anomaly yields a prediction also for  $\Gamma_{\eta' \rightarrow \gamma\gamma}$   
Can use this to pin down all unknowns at NLO

Kaiser 1997



# $\eta$ and $\eta'$ at large $N_c$




Figure taken from diploma work of Roland Kaiser (1997)

Tilted lines: value of  $S = m_s/m_{ud}$ , rectangle: experiment

Central value found in this determination:  $S = 26.6$

Barely differs from leading order result:  $S = 25.9$